

ALL POWER LABS Moving Carbon from Sky to Soil

Agronomic and Carbon Capture Applications of Biochar

By Austin Liu

Austin Liu

Biochar Investigator and Technical Sales at ALL Power Labs

austin@allpowerlabs.com



Table of Contents

- 1. Biochar as carbon storage
- 2. Negative priming
- **3.** Co-composting: the force multiplier
- 4. What gasification does for biochar

Chapter 1: Biochar as carbon storage

Charcoal concepts

Fixed Carbon vs. Volatile carbonReverse-Coal Mining


What is Biochar?

- Fixed Carbon vs volatile carbon
- Reverse coal mining

Fixed Carbon vs. Volatile Carbon

Woody biomass breaks down into two major fractions:

- Fixed carbon, which remains as charcoal
- Volatile carbon, which comes off as smoke (tar gases)



 $\sim 80\%$ of the *mass* of woody biomass consists of volatiles, 20% of the *mass* remains as fixed carbon, between the two, there is roughly 1% ash.

Roughly 50% of the *carbon* is in each fraction.

The fixed carbon fraction is taken out of the carbon cycle as biochar, but the volatile fraction is necessarily burned.

What is Biochar?

- Fixed Carbon vs volatile carbon
- Reverse coal mining

Fixed Carbon vs. Volatile Carbon

50% of the carbon embodied as fixed carbon gets taken out of the carbon cycle as biochar, but the volatile fraction is burned to provide heat for pyrolysis.

- Since the volatile carbon came from the atmosphere to begin with, and since biomass would normally decay back into CO₂, the emissions are carbon neutral in the time frame of decay.
- But the fraction that is removed from the carbon cycle makes this process net carbon negative.

What is Biochar?

- Fixed Carbon vs volatile carbon
- Reverse coal mining

Reverse Coal Mining

The trajectory of carbon in biochar amounts to:

- Capturing carbon out of the air with plants
- Making black carbon out of the plant biomass
 - \circ $\;$ Optionally generating electricity and useful heat
- Burying it in the ground

This is essentially *reverse coal mining*. Biochar sequesters *carbon*, not carbon dioxide.

- 1 ton of carbon is equivalent to 3.7 tons of carbon dioxide. Most of the mass of CO_2 comes from the oxygen.
- Sequestered CO₂ takes two oxygen atoms per carbon with it, but C does not.

Advantages of biochar for carbon storage

- It is *permanent*
- It is *precise*
- It is *productive*
- It is *practical*

Advantages of biochar for carbon storage

- It is *permanent*
- It is precise
- It is *productive*
- It is *practical*

bit.ly/PermanentBiochar





Biochar will never decompose to CO_2 if it is exposed to temperatures high enough to transform its microstructure into a form that is indigestible to microbes. Microbial enzymes can't break down graphitic carbon.




Figure 1: Schematic representation of different molecular forms of carbon in biochar.

Advantages of biochar for carbon storage

• It is *permanent*


- It is *precise*
- It is *productive*
- It is *practical*

bit.ly/PermanentBiochar





Published works on the permanence of biochar



*Fig. 2: Sequestration curve of a 1000 tons carbon sink made from soil-applied biochar with an H:C ratio below 0.4. The persistent aromatic carbon (PAC) pool presents 750 t carbon that will remain over more than 1000 years in the terrestrial system. The semi-persistent carbon (SPC) pool has a minimum MRT of 50 years and was modeled on a bi-modal exponential decay function. The complete SPC decay occurs over 350 years. Thus, the total carbon sink decreases to 87.5% after 50 years and reaches the stable PAC plateau of 75.0% of total pyrogenic carbon after 350 years. The decay function is <i>[Total PyC] ((x))=a*e^((-kf*x))+b*e^((-ks*x))+P*

with a = 45.423, kf = 0.513, b = 212.007, ks = 0.009448, P = 742.5 nd x = year after soil application. The decay curve of the semi-persistent carbon pool is an approximation covering multiple discrete (physical) degradation events rather than a continually harmonious decomposition.

Advantages of biochar for carbon storage

- It is *permanent*
- It is *precise*
- It is *productive*
- It is *practical*

Precision of Biochar Carbon Accounting

Biochar affords a measure of precision and confidence to carbon accounting that is difficult if not impossible for other methods of carbon storage:

- Biochar can be lab tested to determine its precise carbon percentage
- Biochar can be weighed to calculate the precise carbon content
- Biochar can be traded like a commodity

In contrast, accounting for carbon drawdown in living forests, rangelands, and in enhanced weathering schemes is difficult and imprecise.

For the purposes of carbon markets, the precision afforded by biochar is a huge advantage.

Advantages of biochar for carbon storage

- It is *permanent*
- It is *precise*
- It is *productive*
- It is *practical*

Productivity of Biochar Carbon

Other modes of carbon storage are not productive; they passively store carbon. In contrast, the carbon in biochar is *productive*:

- Biochar can radically enhance plant productivity.
- Biochar can trigger forces of nature that store additional carbon without requiring additional human input.
- Biochar can be co-produced with electricity and heat

Biochar used in agriculture is carbon sequestration that can yield a return on investment. Biochar can carry out long term carbon storage right now, without any contingencies on yet undiscovered scientific breakthroughs.

Capturing carbon out of the atmosphere is tough due to the concentration of $\rm CO_2$ being about 0.039%, but plants already do it for free. We just have to keep the $\rm CO_2$ from coming back out.

Chapter 2: Negative Priming

What is Negative Priming?

And why do we want it?

- What is Negative Priming (NP)?
- The 'Five Spheres'
- NP and the Keeling Curve
- Historic Negative Priming
- Negative Priming with BC, Compost
- Negative Priming with Co-composted biochar

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-composte biochar

What is negative priming?

Negative priming is a phenomenon where the addition of carbon to soil triggers the soil to store more and more carbon without additional intervention. Think of it as a deposit of carbon that earns interest.

- The *negative* in *negative priming* comes from 'carbon negative'—taking carbon out of the atmosphere.
- The *priming* in *negative priming* means a stimulus or an initial action, like priming a pump with the liquid to be pumped.

The priming effect in soil describes the loss of stable carbon from soil after fresh organic matter is added. Negative priming describes the opposite.


- What is negative priming (NP)?
- The 'five spheres
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-composter biochar

bit.ly/microbialCsequestration



What is negative priming?

Negative priming is largely due to stabilized microbial organic matter.



A schematic illustration of the changes in soil carbon fractions when new forest residues are added, taken from Figure 12.5 from the Soil Organic Matter chapter of *The Nature and Properties of Soils*, 15th ed. The priming effect is called out in this figure, but negative priming is not labeled in the original graphic. Negative priming is the subsequent increase in soil humus levels following the reduction in microbial biomass.

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-compostebiochar

The 'five spheres'

Carbon can reside in one of five 'spheres'

- 1. the **atmosphere** air
- 2. the hydrosphere rivers, lakes, and oceans
- 3. the **biosphere** living organisms
- 4. the lithosphere rocks and minerals
- 5. the **pedosphere** —soils


Soil is a mixture of air, water, living organisms, minerals, and the necromass of organism—soil organic matter.

To fight climate change, we want to get carbon out of the atmosphere into the other 'spheres'. *The pedosphere is the only sphere where negative priming occurs.*

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-composte biochar

Negative priming and the Keeling curve

The Keeling Curve is the name of the graph that first showed the alarming rise of CO_2 in the atmosphere. This curve has a saw-tooth shape that is due to seasonal plant growth and decay. Plant growth can sharply draw down atmospheric CO_2 ; decay can sharply increase it.



- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NP
- NP with biochar, compost
- NP with co-composted biochar

bit.ly/PerspectiveOnTerraPreta



Historic negative priming

The most compelling historic example of negative priming is the phenomenon of Terra Preta reproducing itself.



Terra Preta is amazonian dark earth, an ancient biochar soil famous for being permanently fertile, with yields double to triple the yields of the weathered tropical soils around it.

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NP
- NP with biochar, compost
- NP with co-composted biochar

bit.ly/PerspectiveOnTerraPreta



Historic negative priming

Terra preta reproduces itself (minus the charcoal content) from forest litter landing on it, like sourdough starter to which fresh flour is added. It so reliably reproduces itself that locals actually mine it and sell it as compost, leaving it time to grow back..



Terra preta being mined. From the 2011 BBC Documentary on terra preta titled "The Secret Of Eldorado"

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-composted biochar


bit.ly/4dcFDcK



Negative priming with biochar and compost

Biochar has been confirmed to trigger negative priming, enough to compensate for the carbon released from burning the volatile carbon in biomass during its production.





John Wick of the Marin Carbon Project has confirmed that scattering compost on range land can trigger negative priming. After a single application of compost to range land, he has observed soil carbon increasing for 15 years. This was reproduced in a second study in a different climate with a different soil type. Featured in the book *Kiss the Ground*.

See <u>https://marincarbonproject.org/</u> for more information

- What is negative priming (NP)?
- The 'five spheres'
- NP and the Keeling curve
- Historic NI
- NP with biochar, compost
- NP with co-composted biochar

Negative priming with co-composted biochar

In terrestrial ecosystems, plants are the only organisms that fix carbon. Bacteria and fungi cannot do this; they only transform plant-derived carbon into more stable forms.

Over the lifetime of a plant, approximately 70-75% of the carbohydrates produced by the plant are put into the soil as root exudates. A portion of these exudates are metabolized, but a portion contributes to microbial biomass which then results in microbial necromass, which makes up much of the stable soil carbon.

Co-composted biochar can radically increase both the plant drawdown and reduce the decomposition of carbon to $\rm CO_2$.

Chapter 3: Co-composting: the force multiplier

Co-composting: the force multiplier

Why co-composting is a universal best practice

- What biochar does to compost
- What compost does to biochar (BC)
- BC abates N2O and methane emissions
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

- What BC does to compost
- What compost does to B0
- Biochar abates N₂O & CH₂
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

What biochar does to compost

Biochar's primary effects on compost accelerated decomposition and much higher composting temperatures.



- What BC does to compost
- What compost does to BC
- Biochar abates N₂O & CH₂
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

bit.ly/BCnutrex




What compost does to biochar

Composting coats biochar with a spongy coating that is responsible for retaining moisture and for all of the nutrient retention and exchange.

Fig. 5

From: Organic coating on biochar explains its nutrient retention and stimulation of soil fertility



This coating appears to play a role in the negative priming observed in co-composted biochar.

- What BC does to compost
- What compost does to BO
- Biochar abates $N_2O \& CH_4$
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

bit.ly/BCvsN2O



Biochar abates $\mathrm{N}_{2}\mathrm{O}$ and methane emissions

scientific reports

Explore content V About the journal V Publish with us V

nature > scientific reports > articles > article

Article Open access Published: 25 April 2013

Biochar and denitrification in soils: when, how much and why does biochar reduce N_2O emissions?

Maria Luz Cayuela, Miguel Angel Sánchez-Monedero, Asunción Roig, Kelly Hanley, Akio Enders & Johannes Lehmann

Scientific Reports 3, Article number: 1732 (2013) Cite this article

23k Accesses | 486 Citations | 19 Altmetric | Metrics

Abstract

Agricultural soils represent the main source of anthropogenic N₂O emissions. Recently, interactions of black carbon with the nitrogen cycle have been recognized and the use of biochar is being investigated as a means to reduce N₂O emissions. However, the mechanisms of reduction remain unclear. Here we demonstrate the significant impact of biochar on denitrification, with a consistent decrease in N₂O emissions by 10–90% in 14 different agricultural soils. Using the ¹⁵N gas-flux method we observed a consistent reduction of the

Electrically conductive biochar *radically* abates N_2O and methane emissions from compost, and continues to abate N_2O in soil. Emissions abatements of both N_2O and methane have been observed to be over 90%.

- What BC does to compost
- What compost does to BO
- Biochar abates N₂O & CH₄
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

Co-composted biochar and plant fertility

Co-composted biochar enables high levels of plant productivity with minimal fertilizer due to radically enhanced ionic nutrient retention and exchange (CEC and AEC).



- What BC does to compost
- What compost does to BO
- Biochar abates $N_2O \& CH_4$
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

Co-composted biochar and plant fertility

Co-composted biochar enables high levels of plant productivity with minimal fertilizer due to radically enhanced ionic nutrient retention and exchange (CEC and AEC).



- What BC does to compost
- What compost does to BO
- Biochar abates N₂O & CH₄
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

Co-composted biochar and plant fertility

Co-composted biochar enables high levels of plant productivity with minimal fertilizer due to radically enhanced ionic nutrient retention and exchange (CEC and AEC).



- What BC does to compost
- What compost does to B
- Biochar abates N₂O & CH₂
- Co-composted BC and plant fertility
- Co-composed BC and negative priming

Co-composted biochar and negative priming

Co-composed biochar enhances negative priming at several levels:

- Adding biochar (black carbon) appears to bias compost to make stable solids rather than CO₂
- Boosting plant productivity boosts root exudates (green carbon), which feeds the soil biome
- Co-composted biochar stimulates soil (and therefore carbon stabilization) by fungi (white carbon).




Chapter 4: What gasification does for biochar

Unexpected and unforeseen benefits

- The five processes of gasification
- Reduction reactions
- Four consequences
- The gasification trifecta

- The five processes
- Reduction reactions
- Four consequences
- The gasification trifecta




* tar cracking is the breakdown of tar into H2, CO, and other flammable gases by exposure to high temperatures.

- The five processes
- The reduction reactions
- Four consequences
- The gasification trifecta

The reduction reactions

(We mean reduction as in the opposite of oxidation.)




- The five processes
- Reduction reactions
- The four consequences
- The gasification trifecta

The four consequences of gasification on BC

Making biochar via gasification has four major consequences:

- 1. Conductivity
- 2. Recalcitrance
- 3. Cleanliness
- 4. Activation



- The five processes
- Reduction reactions
- The four consequences
- The gasification trifecta

bit.ly/conductiveBC



Conductivity is critical

nature communications

Explore content v About the journal v Publish with us v

nature > nature communications > articles > article

Article Open access Published: 31 March 2017

Rapid electron transfer by the carbon matrix in natural pyrogenic carbon

<u>Tianran Sun</u>[™], <u>Barnaby D. A. Levin, Juan J. L. Guzman, Akio Enders, David A. Muller, Largus T.</u> <u>Angenent & Johannes Lehmann</u>

Nature Communications 8, Article number: 14873 (2017) Cite this article

15k Accesses | 385 Citations | 51 Altmetric | Metrics

Abstract

Surface functional groups constitute major electroactive components in pyrogenic carbon. However, the electrochemical properties of pyrogenic carbon matrices and the kinetic preference of functional groups or carbon matrices for electron transfer remain unknown. Here we show that environmentally relevant pyrogenic carbon with average H/C and O/C ratios of less than 0.35 and 0.09 can directly transfer electrons more than three times faster than the charging and discharging cycles of surface functional groups and have a 1.5 V

- The five processes
- Reduction reactions
- The four consequences
- The gasification trifecta

bit.ly/PAHsinBC





Cracking and char cleanliness



Environment International Volume 121, Part 1, December 2018, Pages 169-177



ENTHALPY ANALYTICAL

Analysis Results for 490130

Sample ID: PP30 V3.0 PROTO 1 PISTACHIO SHELL						490130-002 Miscell.	Collected: 08/03/23		
490130-002 Analyte	Result	Qual	Units	RL	DF	Batch	Prepared	Analyzed	Chemis
Method: EPA 8270C-SIM Prep Method: EPA 3546	0	>							
1-Methylnaphthalene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	WLT
2-Methylnaphthalene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Naphthalene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Acenaphthylene	ND	1	ug/Kg	4,700	470	320092	06/10/23	08/11/23	TJW
Acenaphthene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Fluorene	ND	1	ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Phenanthrene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Anthracene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Fluoranthene	ND	1	ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Pyrene	ND	1	ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Benzo(a)anthracene	ND	1	ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Chrysene	ND	1	ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Benzo(b)fluoranthene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Benzo(k)fluoranthene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJIW
Benzo(a)pyrene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Indeno(1,2,3-cd)pyrene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Dibenz(a,h)arithracene	ND		ug/Kg	4,700	470	320092	08/10/23	08/11/23	TJW
Benzo(g,h.i)perylene	ND		ug/Kg	4,700	470	320092	06/10/23	08/11/23	TJW
Surrogates	0		1000	Limits					
Nitrobenzene-d5	58%		%REC	27-125	470	320092	08/10/23	08/11/23	TJJW
2-Fluorobiphenyl	62%		%REC	30-120	470	320092	08/10/23	08/11/23	TJNV
Terphonyl-d14	36%		%REC	33-155	470	320092	08/10/23	08/11/23	TJW

ND Not Detected

Application of biochar to soils may result in plant contamination and human cancer risk due to exposure of polycyclic aromatic hydrocarbons

Jian Wang,^a, <u>Kang Xia</u>^b, <u>Michael Gatheru Waigi</u>^a, <u>Yanzheng Gao</u>^a, <u>X</u> 🖾 , <u>Emmanuel Stephen Odinga</u>^a, <u>Wanting Ling</u>^a, <u>Juan Liu</u>^a

Show more 🗸

+ Add to Mendeley	🔩 Share	55 Cite	
ttps://doi.org/10.1016/j.	anvint 2018 09	1010 7	

Under a Creative Commons license 🛪

Get rights and content 7

open access


Highlights

- The PAH concentrations in biochars were $638\text{--}12,347\,\mu\text{g}/\text{kg}$ depending on production parameters.
- · Root exudates of vegetables enhanced the PAH release from biochars.

- The five processes
- Reduction reactions
- The four consequences
- The gasification trifecta

Reduction and activation

Reduction reactions perforate the char; reduction reactions are what 'activate' activated carbon.



Without reduction

With reduction

- The five processes
- Reduction reactions
- Four consequences
- The gasification trifecta

The gasification trifecta

Biochar made through gasification is what enables "reverse coal mining". The gasification-based biochar trifecta is as follows:

- On-demand renewable energy
- Regenerative agriculture
- Carbon sequestration with benefits

Introducing the machine that does all this...